



AHA 2025 VESALIUS-CV Primary Results Presentation Slides

AMGEN

The Effect of EVolocumab in PatiEntS at High CArdiovascuLar Risk WithoUt Prior Myocardial Infarction or Stroke: Primary Results of the VESALIUS-CV Study

ANDREAE VESALII
BRUXELLENSIS, SCHOLAE

Erin Bohula, MD DPhil

On behalf of the VESALIUS-CV Investigators

*American Heart Association Scientific Sessions
November 8, 2025*

- **LDL-C is a well-established, modifiable CV risk factor**
- **Lowering LDL-C with PCSK9 inhibitors, including evolocumab, ↓ the risk of CV events in patients with a prior, major ASCVD event, such as MI or stroke**
- **Clinical benefit of PCSK9 inhibition in patients without a prior MI or stroke is unknown**

No Known ASCVD

Known ASCVD

Increasing risk for development of ASCVD

Increasing severity of ASCVD

Low-CV Risk
(w/o known ASCVD)

High-CV Risk
(w/o known ASCVD)

Atherosclerosis
w/o Prior Event

Major ASCVD
Event

Diabetes
+ risk enhancer

Coronary disease,
Cerebrovascular disease, or
Peripheral artery disease,
+ risk enhancer
No MI or Stroke

Prior MI
Prior Stroke
Symptomatic PAD
+ risk enhancer

VESALIUS-CV

FOURIER and
ODYSSEY-OUTCOMES

N= 12,257

Event & f/up driven trial:

- 3-P MACE ≥ 751 events
- 4-P MACE $\geq 1,254$ events
- Median f/up ≥ 4.5 yrs

**Stable patients at high-risk for CV events
but no prior MI or stroke***

**LDL-C ≥ 90 mg/dL or
non-HDL-C ≥ 120 mg/dL or
ApoB ≥ 80 mg/dL**

On optimized statin therapy ($\pm$ ezetimibe)

***At least one of the following:**

- CAD without MI
- CVD without stroke
- PAD
- High-risk diabetes mellitus

**Evolocumab SC
140 mg Q2W**

**1:1
Randomization**

**Placebo SC
Q2W**

Dual Primary Endpoints:

Time to coronary heart disease death, MI, or ischemic stroke (3-P MACE)
Time to 3-P MACE plus ischemia-driven arterial revascularization (4-P MACE)

TIMI Study Group

Marc Sabatine (Study Chair)
Robert Giugliano (Co-Principal Investigator)
Erin Bohula (Co-Principal Investigator)
Nicholas Marston (Investigator)

P. Fish, A. Pricken, R. Striar, S. Ewing (Ops)
S. Murphy, J. Kuder, JG. Park (Stats)
Christian Ruff (CEC)

Amgen (Sponsor)

Narimon Honarpour
E. Magnus Ohman
Gabriel Paiva da Silva Lima

Ajay Bhatia, Marcoli Cyrille, Vineet Shastri
J. Bayley, G. Harding, Y. Potapova (Ops)
H. Wang, L. Liu, E. Walsh (Stats)

Executive Committee

Marc Sabatine (Chair)
Robert Giugliano
Jose Nicolau

Lawrence Leiter
Gaetano De Ferrari

Independent Data Monitoring Committee

Charles Hennekens (Chair)
Felicita Andreotti

W. Virgil Brown
Barry Davis

774 enrolling sites from 33 countries




- 12,257[†] in analytic population
- Over a median follow up of 4.6 years, annualized rates of:
 - Premature drug d/c: 4.5%
 - Withdrawal of consent: 0.3%
 - Lost to follow-up: 0.08%
- Follow up for primary end points for 98.7% of total potential patient-years

Demographics		N=12,257	Lipid-lowering therapy (LLT)	
Age (years)		66 [60, 71]	Any LLT	92%
Female		43%	High-intensity LLT regimen	72%
White		93%	Any statin	87%
Hispanic		17%	High-intensity statin	68%
Diabetes		58%	Ezetimibe	20%
Qualifying Disease Categories*			Lipid Values (mg/dL)	
Any qualifying atherosclerosis		67%	LDL-C	122 [104, 149]
CAD w/o MI		45%	Non-HDL-C	153 [130, 182]
CVD w/o stroke		10%	Apolipoprotein-B	100 [89, 121]
PAD		17%		
High-risk diabetes		49%		
With no qualifying atherosclerosis		33%		


% or median and interquartile range.

Pooled data; no difference between treatment arms.


*Qualifying disease categories of CAD CVD, PAD and high-risk diabetes are not mutually exclusive



3-P MACE (CHD death, MI, or ischemic stroke)



4-P MACE (CHD death, MI, ischemic stroke, or IDR)



Dual Primary Endpoints

3-P MACE



HR (95% CI)

0.75 (0.65, 0.86)

p-value

<0.0001

4-P MACE



0.81 (0.73, 0.89)

<0.0001

Key Secondary Endpoints (in testing order)

MI, ischemic stroke, or IDR



0.79 (0.72, 0.88)

<0.0001

Composite
Endpoints

CHD death, MI, or IDR



0.79 (0.72, 0.88)

<0.0001

CV death, MI, or ischemic stroke



0.73 (0.64, 0.84)

<0.0001

CHD death or MI



0.73 (0.62, 0.87)

0.0003

MI



0.64 (0.52, 0.79)

<0.0001

IDR



0.79 (0.70, 0.88)

<0.0001

Individual
Endpoints

CHD death



0.89 (0.68, 1.16)

0.39

CV death



0.79 (0.64, 0.98)

0.032*

All-cause death



0.80 (0.70, 0.91)


0.0005*

Ischemic stroke




0.79 (0.62, 1.01)


0.062*




*Nominal p-value

CHD, coronary heart disease
IDR, ischemia-driven arterial revasc






Major Coronary Event [MCE]
(Death due to AMI, MI)



Major Vascular Event [MVE]
(MCE, fatal/non-fatal stroke, cor revasc)



In pts at high CV risk w/o prior MI or stroke, addition of the PCSK9 inhibitor, evolocumab, to baseline LLRx resulted in:

- Median achieved LDL-C of 45 mg/dL (1.16 mmol/L)
- 25% ↓ in 3-P MACE and 19% ↓ in 4-P MACE
- Consistency across subgroups, including in those with diabetes and no qualifying atherosclerosis
- 27% ↓ in CV death, MI or ischemic stroke and 36% ↓ in MI
- Nominally lower rates of CV and all-cause death

Guidelines have progressively recommended lower LDL-C goals of <70 or <55 mg/dL in very high-risk patients, and most recently <40 mg/dL in extreme-risk patients.

The reduction in MACE seen in VESALIUS-CV, with an achieved LDL-C of ~40 mg/dL in the evolocumab arm, supports intensive LDL-C lowering to this level, even in patients without a prior major ASCVD event.



ORIGINAL ARTICLE

Evolocumab in Patients without Previous
Myocardial Infarction or Stroke

Erin A. Bohula, M.D., D.Phil.,¹ Nicholas A. Marston, M.D., M.P.H.,¹
Ajay K. Bhatia, M.D., Ph.D.,² Gaetano M. De Ferrari, M.D.,³
Lawrence A. Leiter, M.D.,⁴ Jose C. Nicolau, M.D.,⁵ Jeong-Gun Park, Ph.D.,¹
Julia F. Kuder, M.A.,¹ Sabina A. Murphy, M.P.H.,¹ Emileigh Walsh, Ph.D.,²
Huei Wang, Ph.D.,² Vladimir Blaha, M.D., Ph.D.,⁶ Andrzej Budaj, M.D., Ph.D.,⁷
Jan H. Cornel, M.D., Ph.D.,⁸ Assen Goudev, M.D.,⁹ Robert Gabor Kiss, M.D.,¹⁰
Alberto J. Lorenzatti, M.D.,¹¹ Alexander Parkhomenko, M.D.,¹²
Marcoli Cyrille, M.D.,² Gabriel Paiva da Silva Lima, M.D.,²
E. Magnus Ohman, M.D.,² Robert P. Giugliano, M.D.,¹ and
Marc S. Sabatine, M.D., M.P.H.,¹ for the VESALIUS-CV Investigators*

